

UCLA

UCLA Previously Published Works

Title

Influences of light and humidity on carbonyl sulfide-based estimates of photosynthesis.

Permalink

<https://escholarship.org/uc/item/7vt4f657>

Journal

Proceedings of the National Academy of Sciences of the United States of America,
116(7)

ISSN

0027-8424

Authors

Kooijmans, Linda MJ
Sun, Wu
Aalto, Juho
[et al.](#)

Publication Date

2019-02-01

DOI

10.1073/pnas.1807600116

Peer reviewed

Influences of light and humidity on carbonyl sulfide-based estimates of photosynthesis

Linda M. J. Kooijmans^{a,1,2}, Wu Sun^b, Juho Aalto^{c,d}, Kukka-Maaria Erkkilä^c, Kadmiel Maseyk^e, Ulrike Seibt^b, Timo Vesala^{c,f}, Ivan Mammarella^c, and Huilin Chen^{a,2}

^aCentre for Isotope Research, University of Groningen, 9747 AG, Groningen, The Netherlands; ^bDepartment of Atmospheric and Oceanic Sciences, University of California, Los Angeles, CA 90095-1565; ^cFaculty of Science, Institute for Atmospheric and Earth System Research/Physics, University of Helsinki, 00014 Helsinki, Finland; ^dStation for Measuring Forest Ecosystem–Atmosphere Relations II, Hyytiälä Forestry Field Station, University of Helsinki, 35500 Korkeakoski, Finland; ^eSchool of Environment, Earth and Ecosystem Sciences, The Open University, MK 7 6AA Milton Keynes, United Kingdom; and ^fFaculty of Agriculture and Forestry, Institute for Atmospheric and Earth System Research/Forest Sciences, University of Helsinki, 00014 Helsinki, Finland

Edited by Steven C. Wofsy, Harvard University, Cambridge, MA, and approved December 18, 2018 (received for review May 2, 2018)

Understanding climate controls on gross primary productivity (GPP) is crucial for accurate projections of the future land carbon cycle. Major uncertainties exist due to the challenge in separating GPP and respiration from observations of the carbon dioxide (CO₂) flux. Carbonyl sulfide (COS) has a dominant vegetative sink, and plant COS uptake is used to infer GPP through the leaf relative uptake (LRU) ratio of COS to CO₂ fluxes. However, little is known about variations of LRU under changing environmental conditions and in different phenological stages. We present COS and CO₂ fluxes and LRU of Scots pine branches measured in a boreal forest in Finland during the spring recovery and summer. We find that the diurnal dynamics of COS uptake is mainly controlled by stomatal conductance, but the leaf internal conductance could significantly limit the COS uptake during the daytime and early in the season. LRU varies with light due to the differential light responses of COS and CO₂ uptake, and with vapor pressure deficit (VPD) in the peak growing season, indicating a humidity-induced stomatal control. Our COS-based GPP estimates show that it is essential to incorporate the variability of LRU with environmental variables for accurate estimation of GPP on ecosystem, regional, and global scales.

carbonyl sulfide | photosynthesis | stomatal conductance | carbon cycle

Carbonyl sulfide (COS) follows the same diffusion pathway into the leaf chloroplasts as CO₂ and is consumed by the enzyme carbonic anhydrase (CA) (1, 2). The hydrolysis of COS via CA is irreversible (3), such that no respiration-like COS flux is evident under ambient conditions. Consequently, the atmospheric drawdown of COS above an ecosystem reflects the uptake of COS by plants, provided that other sources and sinks in the ecosystem are negligible or known. The dominant vegetative sink of COS was therefore recognized as a way to separate net ecosystem exchange of CO₂ (NEE) into gross primary productivity (GPP) and respiration (4–7). With a known ratio of COS to CO₂ uptake at the leaf level, GPP can be determined from COS ecosystem fluxes (F_{COS-E}) following (5, 7):

$$GPP_{COS} = -F_{COS-E} \frac{C_{a,COS}}{C_{a,CO_2}} \frac{1}{LRU}, \quad [1]$$

with atmospheric mole fractions $C_{a,COS}$ and C_{a,CO_2} , and the leaf-scale relative uptake ratio (LRU) = $F_{COS}/F_{CO_2} \cdot C_{a,CO_2}/C_{a,COS}$, with F_{COS} and F_{CO_2} being the flux rates of COS and CO₂ at the leaf level. LRU is also referred to as the ratio of deposition velocities of COS and CO₂ (8). The accuracy of LRU is key in translating COS fluxes into GPP, and several studies have derived LRU for different plant species from chamber enclosure measurements (8–17). Those LRU values ranged from 0.4 to 9.5 with a median of 1.75 and with 50% of the values between 1.48 and 2.46 around the median (see ref. 18 for an overview).

Many of the laboratory studies measured LRU under constant conditions and few have investigated LRU response to environmental variations or under field conditions (13, 17). If effects of light, humidity, and temperature on dissolution, diffusion, and relevant enzyme reactions differ between COS and CO₂, then LRU should be expected to vary (13). It has already been found that LRU changes with light intensity (13, 14, 17, 19, 20). This is due to the light independence of the CA enzyme that controls F_{COS} (14, 21, 22), whereas F_{CO_2} depends on the light reactions in the photosystems.

LRU values are typically larger than 1.0, which implies that the deposition velocities of COS are typically higher than those of CO₂. This is attributed to a lower reaction efficiency of ribulose-1,5-bisphosphate carboxylase/oxygenase with CO₂ than that of CA with COS (9, 12), which can be expected because CA is known to be the enzyme with the highest molar activity (2).

Significance

Carbonyl sulfide (COS) measurements enable quantification of terrestrial photosynthesis, which cannot be directly measured at scales greater than the leaf level. The accuracy of COS-based estimates of gross primary production (GPP) depends on how we relate the COS uptake to that of CO₂. This study shows that COS-based GPP estimates will be significantly overestimated if the different environmental responses of COS and CO₂ uptake are not taken into account. These findings are relevant for studies that rely on COS to quantify ecosystem to regional scale GPP, and support the use of a COS-based approach to constrain ecosystem flux partitioning. Moreover, the strong stomatal control on COS uptake shown in this study makes COS a suitable tracer for stomatal diffusion.

Author contributions: L.M.J.K., W.S., J.A., K.-M.E., K.M., U.S., T.V., I.M., and H.C. designed research; L.M.J.K., W.S., J.A., K.-M.E., K.M., and H.C. performed research; L.M.J.K., W.S., J.A., K.-M.E., and H.C. analyzed data; and L.M.J.K., W.S., J.A., K.-M.E., K.M., U.S., T.V., I.M., and H.C. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

This open access article is distributed under Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND).

Data deposition: The data used in this work are available from <https://zenodo.org/record/1211481#XB4Lb9IzblU>. The dataset includes branch fluxes and mole fractions of COS and CO₂, LRU, stomatal conductance, internal conductance, eddy-covariance fluxes of COS and CO₂, GPP, and meteorological information. The code used to calculate the chamber fluxes from measured mole fractions is available from <https://zenodo.org/record/1197330#XB4OLNlZblU>. The code used to generate the main results in this work is available from <https://zenodo.org/record/1211499#XB4PStIzblU>.

¹Present address: Meteorology and Air Quality Group, Wageningen University and Research Centre 6700 AA, Wageningen, The Netherlands.

²To whom correspondence may be addressed. Email: linda.kooijmans@wur.nl or huiLin.chen@rug.nl.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1807600116/-DCSupplemental.

Published online January 25, 2019.




Fig. 2. Responses of F_{cos} , F_{co_2} , and LRU to light and of F_{cos} to $g_{\text{s,cos}}$. Average F_{cos} , F_{co_2} (A and B) and LRU (C and D) versus PAR, and F_{cos} versus $g_{\text{s,cos}}$ (E and F) from 18 May to 13 July for chambers 1 (Left) and 2 (Right). Data are plotted as the median of 15 equal-sized bins in the x range. The error bars represent the 25th and 75th percentiles of data in each bin. For the correlation of F_{cos} with $g_{\text{s,cos}}$ (E and F) the different colors represent different light conditions: nighttime (blue); daytime with low light conditions (PAR < 150 and 100 $\mu\text{mol m}^{-2}\text{s}^{-1}$ for chambers 1 and 2, respectively; green); daytime with high light conditions (PAR > 300 $\mu\text{mol m}^{-2}\text{s}^{-1}$; orange). A transition phase between low and high PAR values is neglected. The coefficient of determination (R^2), slope (sl), significance level (P), and number of data (n) are given for a linear regression through the median values (E and F).

example, Kesselmeier and Merk (9) determined LRU at a light level of 300 $\mu\text{mol m}^{-2}\text{s}^{-1}$ and Sandoval-Soto et al. (8) also measured LRU in Scots pine but at a light level of 600 $\mu\text{mol m}^{-2}\text{s}^{-1}$ where F_{co_2} is not PAR saturated.

Internal Conductance of COS Limits F_{cos} During Daytime. We estimated the internal conductance to COS ($g_{\text{i,cos}}$), which is a combination of nonstomatal conductance terms, and find that during daytime $g_{\text{i,cos}}$ is smaller than $g_{\text{s,cos}}$ (see SI Appendix, Fig. S4 and the accompanying explanation). The ratio of $g_{\text{s,cos}}$ over $g_{\text{i,cos}}$ determines the relative importance of the two conductances on F_{cos} and thereby also on LRU (see equation 8 in ref. 12). The fact that we find a relatively low $g_{\text{i,cos}}$ compared with $g_{\text{s,cos}}$ during the daytime implies that $g_{\text{i,cos}}$ has a relatively large control on F_{cos} . Wehr et al. (23) estimated that the biochemical conductance (the CA activity) was of similar magnitude as $g_{\text{s,cos}}$ during the daytime. The fact that we also find a relatively high importance of $g_{\text{i,cos}}$ emphasizes the need to take into account $g_{\text{i,cos}}$ on the total conductance of COS uptake (17). The day–night difference of $g_{\text{s,cos}}$ is larger than that of $g_{\text{i,cos}}$, and therefore $g_{\text{s,cos}}$ has a relatively larger effect on day–night differences of F_{cos} than $g_{\text{i,cos}}$ has. This means that the diurnal change of F_{cos} is largely controlled by $g_{\text{s,cos}}$. Furthermore, $g_{\text{i,cos}}$ has a relatively larger limiting role on F_{cos} during daytime than during nighttime (SI Appendix, Fig. S4). This variable role of $g_{\text{i,cos}}$ over a day explains why the relation between F_{cos} and $g_{\text{s,cos}}$ is different between different moments of the day, as depicted by different light levels in Fig. 2 E and F. If F_{cos} is used to determine $g_{\text{s,cos}}$,

and the limiting role of $g_{\text{i,cos}}$ on F_{cos} is ignored, this would lead to underestimation of daytime $g_{\text{s,cos}}$. When the $F_{\text{cos}}-g_{\text{s,cos}}$ relationship is assumed to be the same for daytime and nighttime (following the blue curve in Fig. 2 E and F), $g_{\text{s,cos}}$ would be equal to 0.012 and 0.020 $\text{mol m}^{-2}\text{s}^{-1}$ for chambers 1 and 2, respectively, at F_{cos} of $-3 \text{ pmol m}^{-2}\text{s}^{-1}$ (the average F_{cos} at high light levels). These values are, respectively, 46% and 48% smaller than what is actually observed (following the orange curve in Fig. 2 E and F). Therefore, ignoring the role of $g_{\text{i,cos}}$ would lead to a substantial underestimation of $g_{\text{s,cos}}$.

Seasonal Variation of LRU Influenced by Environmental Variables.

Fig. 3 shows the light-saturated LRU per month binned by VPD. The monthly median LRU decreases by 0.2 from April to July. No significant correlation between LRU and VPD can be detected before June, whereas a significant decrease of LRU with VPD is observed in June and July (indicated by the significance levels in Fig. 3). The fact that the LRU–VPD correlation follows the progression of the growing season is associated with the increase of daytime VPD. Early in the season F_{cos} and F_{co_2} are not solely limited by stomatal conductance but rather by low temperatures, as is shown in SI Appendix, Fig. S5. The low temperatures suppress enzyme activities or mesophyll diffusion and therefore $g_{\text{i,cos}}$ has a relatively larger limiting effect on F_{cos} than $g_{\text{s,cos}}$ early in the season. In the course of the season the limitation of VPD on stomatal conductance becomes stronger, which manifests in the LRU–VPD relationship. This emphasizes that the LRU– $g_{\text{s,cos}}$ correlation (SI Appendix, Fig. S3) only applies when both F_{cos} and F_{co_2} are controlled by stomatal conductance; i.e., at high temperatures and high light conditions.

Light and Humidity-Dependent LRU Required for Accurate COS-Based GPP Estimates.

In Fig. 4 we compare COS-based GPP estimates (GPP_{COS}) from COS ecosystem fluxes (determined from eddy-covariance measurements and subtracted estimates of the soil flux) with GPP from a traditional flux-partitioning method based on extrapolating nighttime respiration to the daytime (28) (GPP_{NEE}). GPP_{COS} is determined using different parameterizations of LRU: (i) a fit of the measured LRU (averaged over chambers 1 and 2) against PAR, which captures the decrease of LRU toward simultaneously increasing VPD and PAR ($\text{GPP}_{\text{COS-fit}}$; see SI Appendix, Fig. S6 for the LRU–PAR relationship) and (ii) LRU fixed at 1.1 (the average LRU that we find at high light levels) and 1.6 [similar to what has been frequently used in other literature (7, 15, 29)], where the latter is shown in Fig. 4 as $\text{GPP}_{\text{COS-const}}$. The shading of the GPP estimates represents the uncertainty based on Monte Carlo sampling of all parameters contributing to the GPP calculations (Methods). The GPP_{COS} uncertainty is larger than that of GPP_{NEE} , partly because the relative uncertainty of COS mole fraction measurements ($\sim 1.7\%$ of a typical ambient level of 450 ppt) is greater than that of CO_2 mole fraction measurements ($\sim 0.06\%$ of a typical value of 400 ppm) (30). Still, Fig. 4 shows that the accuracy of GPP_{COS} is sufficient to detect differences between GPP_{COS} and GPP_{NEE} . We also calculated GPP_{COS} with the measured hourly LRU to determine to what extent uncertainty in the LRU–PAR function adds uncertainty to $\text{GPP}_{\text{COS-fit}}$. The uncertainties did not decrease with measured LRU values compared with the LRU–PAR function, implying that the empirical function captures the variability of LRU over the measurement period well.

With the constant LRU, the earlier peak of F_{cos} leads to an earlier peak in $\text{GPP}_{\text{COS-const}}$ compared with GPP_{NEE} . The peak of ecosystem F_{cos} , and thus that of $\text{GPP}_{\text{COS-const}}$, is 2 h later than the peak of F_{cos} measured at the branch level at the top of the canopy. The reason for the delay between the F_{cos} peak from branches and ecosystem is that the diurnal pattern of the bulk canopy conductance is more symmetric, because light rather than g_{s} is limiting CO_2 assimilation in the lower canopy, in contrast to the top of the canopy (31). When GPP_{COS} is calculated with the average LRU that we find at high light levels (1.1), we find GPP_{COS} ($13.4 \pm 1.3 \text{ g C m}^{-2}\text{d}^{-1}$; daytime data only) to

soil flux of $-2.7 \text{ pmol m}^{-2} \text{ s}^{-1}$ was subtracted from the ecosystem fluxes such that the remaining flux represents the vegetative COS exchange. The averages and uncertainties shown in Fig. 4 are based on 1,000 subsamples of Monte Carlo simulations that include uncertainties of all contributing components in the GPP calculation. That is, the SE of $F_{\text{COS-EC}}$ and NEE; the COS soil flux uncertainty of $1.1 \text{ pmol m}^{-2} \text{ s}^{-1}$ (45); the SE of the fitting parameters of the LRU-PAR relation (using the median PAR in the calculation), or no uncertainty in LRU in the case of a constant LRU; the uncertainties of COS and CO_2 mole fractions of 6.0 ppt and 0.13 ppm respectively (30), and the range of respiration calculations from figure 11 in ref. 47.

Meteorological Data. In addition to the temperature and PAR sensors installed at the branch chambers we use the data that are made available through the SmartSMEAR database that contains continuous data records from all SMEAR sites (available at <https://avaa.tdata.fi/>).

Statistical Tests. The significance of correlations is tested with two-sided t tests of which the significance levels (P) are reported. To reduce the effect

of outliers we test the linear correlation of data based on bin-averaged medians, where the bins are of equal size. The number of samples of the original data are reported for each t test. The number of bins are mentioned in figure legends.

Code and Data Availability. Data and code related to this paper are available in refs. 48–50.

ACKNOWLEDGMENTS. We thank the technical staff at the SMEAR II station in Hyttälä and M. de Vries, B. A. M. Kers, H. A. Been, and H. G. Jansen from the University of Groningen for their help during preparation and maintenance of the field campaign. This research was supported by the Startup Grant (awarded to H.C.) at the University of Groningen, the National Oceanic and Atmospheric Administration Climate Program Office Grant (NA13OAR4310082), the European Union's Horizon 2020 research and innovation program (Grant 654182), the Vilho, Yrjö, and Kalle Väisälä Foundation, Integrated Carbon Observation System-Finland (Grant 281255), Academy of Finland Center of Excellence program (307331), and the NSF CAREER Award 1455381 (to U.S.).

- Protoschill-Krebs G, Wilhelm C, Kesselmeier J (1992) Enzymatic pathways for the consumption of carbonyl sulphide (COS) by higher plants. *Bot Acta* 105:206–212.
- Protoschill-Krebs G, Wilhelm C, Kesselmeier J (1996) Consumption of carbonyl sulphide (COS) by higher plant carbonic anhydrase (CA). *Atmos Environ* 30:3151–3156.
- Notni J, Schenk S, Protoschill-Krebs G, Kesselmeier J, Anders E (2007) The missing link in COS metabolism: A model study on the reactivation of carbonic anhydrase from its hydrosulfide analogue. *Chembiochem* 8:530–536.
- Montzka SA, et al. (2007) On the global distribution, seasonality, and budget of atmospheric carbonyl sulfide (COS) and some similarities to CO_2 . *J Geophys Res Atmos* 112:D09302.
- Campbell JE, et al. (2008) Photosynthetic control of atmospheric carbonyl sulfide during the growing season. *Science* 322:1085–1088.
- Berry J, et al. (2013) A coupled model of the global cycles of carbonyl sulfide and CO_2 : A possible new window on the carbon cycle. *J Geophys Res Biogeosci* 118:842–852.
- Asaf D, et al. (2013) Ecosystem photosynthesis inferred from measurements of carbonyl sulphide flux. *Nat Geosci* 6:186–190.
- Sandoval-Soto L, et al. (2005) Global uptake of carbonyl sulfide (COS) by terrestrial vegetation: Estimates corrected by deposition velocities normalized to the uptake of carbon dioxide (CO_2). *Biogeosciences* 2:125–132.
- Kesselmeier J, Merk L (1993) Exchange of carbonyl sulfide (COS) between agricultural plants and the atmosphere: Studies on the deposition of COS to peas, corn and rapeseed. *Biogeochemistry* 23:47–59.
- Kesselmeier J, et al. (1993) Reduced sulfur compound exchange between the atmosphere and tropical tree species in Southern Cameroon. *Biogeochemistry* 23:23–45.
- Kuhn U, et al. (1999) Carbonyl sulfide exchange on an ecosystem scale: Soil represents a dominant sink for atmospheric COS. *Atmos Environ* 33:995–1008.
- Seibt U, Kesselmeier J, Sandoval-Soto L, Kuhn U, Berry JA (2010) A kinetic analysis of leaf uptake of COS and its relation to transpiration, photosynthesis and carbon isotope fractionation. *Biogeosciences* 7:333–341.
- Stimler K, Montzka SA, Berry JA, Rudich Y, Yakir D (2010) Relationships between carbonyl sulfide (COS) and CO_2 during leaf gas exchange. *New Phytol* 186:869–878.
- Stimler K, Berry JA, Montzka SA, Yakir D (2011) Association between carbonyl sulfide uptake and $(18)\Delta$ during gas exchange in C(3) and C(4) leaves. *Plant Physiol* 157:509–517.
- Stimler K, Berry JA, Yakir D (2012) Effects of carbonyl sulfide and carbonic anhydrase on stomatal conductance. *Plant Physiol* 158:524–530.
- Berkelhammer M, et al. (2014) Constraining surface carbon fluxes using in situ measurements of carbonyl sulfide and carbon dioxide. *Global Biogeochem Cycles* 28:161–179.
- Sun W, Maseyk K, Lett C, Seibt U (2018) Stomatal control of leaf fluxes of carbonyl sulfide and CO_2 in a Typha freshwater marsh. *Biogeosciences* 15:3277–3291.
- Whelan ME, et al. (2018) Reviews and syntheses: Carbonyl sulfide as a multi-scale tracer for carbon and water cycles. *Biogeosciences* 15:3625–3657.
- Maseyk K, et al. (2014) Sources and sinks of carbonyl sulfide in an agricultural field in the Southern Great Plains. *Proc Natl Acad Sci USA* 111:9064–9069.
- Commane R, et al. (2015) Seasonal fluxes of carbonyl sulfide in a midlatitude forest. *Proc Natl Acad Sci USA* 112:14162–14167.
- Gries C, Nash TH, Kesselmeier J (1994) Exchange of reduced sulfur gases between lichens and the atmosphere. *Biogeochemistry* 26:25–39.
- Protoschill-Krebs G, Wilhelm C, Kesselmeier J (1995) Consumption of carbonyl sulphide by chlamydomonas reinhardtii with different activities of carbonic anhydrase (CA) induced by different CO_2 growing regimes. *Bot Acta* 108:445–448.
- Wehr R, et al. (2017) Dynamics of canopy stomatal conductance, transpiration, and evaporation in a temperate deciduous forest, validated by carbonyl sulfide uptake. *Biogeosciences* 14:389–401.
- Geng C, Mu Y (2006) Carbonyl sulfide and dimethyl sulfide exchange between trees and the atmosphere. *Atmos Environ* 40:1373–1383.
- Farquhar GD, von Caemmerer S, Berry JA (1980) A biochemical model of photosynthetic CO_2 assimilation in leaves of C 3 species. *Planta* 149:78–90.
- Cowan IR, Farquhar GD (1977) Stomatal function in relation to leaf metabolism and environment. *Symp Soc Exp Biol* 31:471–505.
- Farquhar GD (1982) Stomatal conductance and photosynthesis. *Annu Rev Plant Physiol* 33:317–345.
- Reichstein M, et al. (2005) On the separation of net ecosystem exchange into assimilation and ecosystem respiration: Review and improved algorithm. *Glob Change Biol* 11:1424–1439.
- Hilton T, et al. (2017) Peak growing season gross uptake of carbon in North America is largest in the Midwest USA. *Nat Clim Chang* 7:450–454.
- Kooijmans LMJ, et al. (2016) Continuous and high-precision atmospheric concentration measurements of COS, CO_2 , CO and H_2O using a quantum cascade laser spectrometer (QCLS). *Atmos Meas Tech* 9:5293–5314.
- Launiainen S, Katul GG, Kolari P, Vesala T, Hari P (2011) Empirical and optimal stomatal controls on leaf and ecosystem level CO_2 and H_2O exchange rates. *Agric For Meteorol* 151:1672–1689.
- Wehr R, et al. (2016) Seasonality of temperate forest photosynthesis and daytime respiration. *Nature* 534:680–683.
- Wilkman E, et al. (2018) Temperature response of respiration across the heterogeneous landscape of the Alaskan Arctic tundra. *J Geophys Res Biogeosci* 123:2287–2302.
- Huang K, et al. (2014) Impacts of diffuse radiation on light use efficiency across terrestrial ecosystems based on eddy covariance observation in China. *PLoS One* 9:e110988.
- Thornton PE, Lamarque J-F, Rosenbloom NA, Mahowald NM (2007) Influence of carbon-nitrogen cycle coupling on land model response to CO_2 fertilization and climate variability. *Global Biogeochem Cycles* 21:1–15.
- Krinner G, et al. (2005) A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system. *Global Biogeochem Cycles* 19:GB1015.
- Sitch S, et al. (2004) Evaluation of ecosystem dynamics plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model. *Glob Change Biol* 9:161–185.
- Launio T, Peylin P, Belviso S, Poulter B (2015) A new model of the global biogeochemical cycle of carbonyl sulfide—Part 2: Use of carbonyl sulfide to constrain gross primary productivity in current vegetation models. *Atmos Chem Phys* 15:9285–9312.
- Yang F, Qubaja R, Tatarinov F, Rotenberg E, Yakir D (2018) Assessing canopy performance using carbonyl sulfide measurements. *Glob Change Biol* 24:3486–3498.
- Bäck J, et al. (2012) Chemodiversity of a Scots pine stand and implications for terpene air concentrations. *Biogeosciences* 9:689–702.
- Aalto J, et al. (2014) New foliage growth is a significant, unaccounted source for volatiles in boreal evergreen forests. *Biogeosciences* 11:1331–1344.
- Kooijmans LMJ, et al. (2017) Canopy uptake dominates nighttime carbonyl sulphide fluxes in a boreal forest. *Atmos Chem Phys* 17:11453–11465.
- Ball JT, Woodrow IE, Berry JA (1987) A model predicting stomatal conductance and its contribution to the control of photosynthesis under different environmental conditions. *Progress in Photosynthesis Research*, ed Biggens J (Springer, Dordrecht, The Netherlands), Vol 4, pp 221–224.
- Nobel PS (2009) *Physicochemical and Environmental Plant Physiology* (Academic, Oxford).
- Sun W, et al. (2018) Soil fluxes of carbonyl sulfide (COS), carbon monoxide, and carbon dioxide in a boreal forest in Southern Finland. *Atmos Chem Phys* 18:1363–1378.
- Mammarella I, Peltola O, Nordbo A, Järvi L, Rannik Ü (2016) Quantifying the uncertainty of eddy covariance fluxes due to the use of different software packages and combinations of processing steps in two contrasting ecosystems. *Atmos Meas Tech* 9:4915–4933.
- Kolari P, et al. (2009) CO_2 exchange and component CO_2 fluxes of a boreal Scots pine forest. *Boreal Environ Res* 14:761–783.
- Kooijmans LMJ (2018) Code for “Influences of light and humidity on carbonyl sulfide-based estimates of photosynthesis.” Zenodo. Available at <https://zenodo.org/record/1211499#.XEC-zFVKgdV>. Deposited April 3, 2018.
- Kooijmans LMJ, et al. (2018) Dataset for “Influences of light and humidity on carbonyl sulfide-based estimates of photosynthesis.” Zenodo. Available at <https://zenodo.org/record/1211481#.XEC-NVVKhQJ>. Deposited April 3, 2018.
- Sun W (2018) Code for calculation of chamber fluxes: “PyChamberFlux.” Zenodo. Available at <https://zenodo.org/record/1197330#.XEC-i1VKgdV>. Deposited March 13, 2018.